

## IUPAC Task Group on Atmospheric chemical Kinetic Data Evaluation – Data Sheet V.A1.19 H119

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>.

This data sheet last evaluated: April 2010; last change to preferred values: April 2010.

### CH<sub>3</sub>C(O)OH + ice

#### Experimental data

Parameter	Temp./K	Reference	Technique/ Comments
<i>Partition coefficients: <math>K_{inc}</math></i>			
$1.9 \times 10^{-11} \exp(6800/T)$	220-245	Sokolov and Abbatt, 2002	CWFT-MS (a)
320	193	Picaud et al., 2005	CWFT-MS (b)
258	203		
153	213		
73	223		
$2.3 \times 10^{-10} \exp(6530/T)$	197-227	von Hessberg et al., 2008	CWFT-MS (c)
$1.9 \times 10^{-12} \exp(7400/T)$	213-243	Kerbrat et al., 2010	PBFT-CIMS (d)
$5.5 \times 10^{-9} \exp(5703/T)$	218-238	Symington et al., 2010	CWFT-MS (e)

#### Comments

- (a) Ice film made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH<sub>3</sub>C(O)OH to ice at various temperatures was analysed using the Langmuir isotherm. The values for  $K_{inc}$  at individual temperatures given in the Table uses the reported values of  $K_{LangP}(T)$  and  $N_{max}(T)$ . No errors were reported. The temperature dependent expression of  $K_{inc}$  was derived by fitting to these five data points. A value of  $\Delta H_{ads} = -73 \pm 12 \text{ kJ mol}^{-1}$  was reported. CH<sub>3</sub>C(O)OH concentration varied between  $\approx 4 \times 10^{10}$  and  $2 \times 10^{13}$  molecule cm<sup>-3</sup>. with between 2 and 20 % present as dimers.
- (b) Ice film, 30-80  $\mu\text{m}$  thick was made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH<sub>3</sub>C(O)OH to ice at various temperatures was analysed using the BET isotherm to derive a value of  $\Delta H_{ads} = -33.5 \pm 4.2 \text{ kJ mol}^{-1}$ . The parameterised BET isotherms were used to calculate values of  $K_{inc}$  at the four temperatures where reversible uptake was observed. The authors suggest that most of the acetic acid was in the form of dimers in their experiments.
- (c) Ice film made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Values of  $N_{max} = 2.4 \times 10^{14}$  molecule cm<sup>-2</sup> (independent of temperature) and  $\Delta H_{ads} = -55 \pm 9 \text{ kJ mol}^{-1}$  were reported. CH<sub>3</sub>C(O)OH concentration varied between  $3 \times 10^9$  and  $2 \times 10^{11}$  molecule cm<sup>-3</sup> and the fraction of dimers present was calculated to be less than 15 % for all temperatures and concentrations except for 197 K and  $[\text{HC(O)OH}] > 2 \times 10^{10}$  molecule cm<sup>-3</sup>.
- (d) Packed ice bed flow tube at atmospheric pressure. Partition coefficients derived from analysis of breakthrough curves and using the geometric ice surface area.
- (e) Ice film made by freezing distilled water. CH<sub>3</sub>C(O)OH concentration varied between  $\approx 3 \times 10^{10}$  and  $2.6 \times 10^{13}$  molecule cm<sup>-3</sup>. Uptake was found to be reversible and equilibrium surface

coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH<sub>3</sub>C(O)OH to ice at various temperatures was analysed using the Langmuir isotherm and  $N_{\max}$  was found to be  $2.4 \times 10^{14}$  molecule cm<sup>-2</sup>. Linear least squares fit of selected data for concentrations  $< 2.5 \times 10^{12}$  molecule cm<sup>-3</sup> was used to determine the optimum values of  $K_{\text{linC}}$ . A value of  $\Delta H_{\text{ads}} = -49 (\pm 8)$  kJ mol<sup>-1</sup>, was reported.

### Preferred Values

Parameter	Value	T/K
$K_{\text{linC}} / \text{cm}$	$1.5 \times 10^{-14} \exp(8500/T)$	195 - 240
$N_{\max} / \text{molecule cm}^{-2}$	$2.5 \times 10^{14}$	
<i>Reliability</i>		
$\Delta(E/R) / \text{K}$	$\pm 1000$	195 - 240
$\Delta \log N_{\max}$	0.1	

### Comments on Preferred Values


There are five experimental studies of the reversible uptake of CH<sub>3</sub>C(O)OH to pure ice surfaces, the results differing somewhat, possibly due to the presence of varying amounts of the acetic acid dimer. Sokolov et al. (2002), von Hessberg et al. (2007), Kerbrat et al. (2010) and Symington et al. (2010) suggest that the fraction of dimers present in their samples was of the order of percent, whereas Picaud et al. (2005) calculated that almost 100 % of their sample was dimerised. It is not clear if this assumption was partially responsible for the much lower partitioning coefficients found by Picaud et al. The published data of Sokolov et al. (2002), von Hessberg et al. (2007), Kerbrat et al., (2010) and Symington et al., (2010) are in good agreement and were used to derive the preferred expression, which indicates an enthalpy of adsorption of  $70 (\pm 10)$  kJ mol<sup>-1</sup>. The values of  $N_{\max}$  returned by the various studies is variable, probably due to extrapolations from low coverages and also due to lateral interactions at high coverages, which make this parameter generally difficult to access experimentally (Jedlovszky et al., 2006).

Theoretical investigations (Compoint et al. 2002; Picaud et al. 2005) have shown that the hydroxyl and carbonyl groups of acetic acid are bound to two surface water molecules, with the CH<sub>3</sub> group directed away from the ice surface. This would tend to suggest that acetic acid dimers, for which OH and CO bonds are no longer available, would undergo significantly weaker interaction with the ice surface.

### References

- Compoint, M., Toubin, C., Picaud, S., Hoang, P. N. M. and Girardet, C.: Chem. Phys. Lett. 365, 1-7, 2002.
- Kerbrat, M., Huthwelker, T., Bartels-Rausch, T., Gäggeler, H. W., and Ammann, M., Phys. Chem. Chem. Phys., in press, 2010.
- Jedlovszky, P., Partay, L., Hoang, P. N. M., Picaud, S., von Hessberg, P. and Crowley, J. N.: J. Am. Chem. Soc. 128, 15300-15309, 2006.
- Picaud, S., Hoang, P. N. M., Peybernes, N., Le Calve, S. and Mirabel, P.: J. Chem. Phys. 122, article 194707, 2005.
- Sokolov, O. and Abbatt, J. P. D.: J. Phys. Chem. 106, 775-782, 2002.
- Symington, A., Cox, R. A. and Fernandez, M. A.: Z. Phys. Chem., in press 2010.

von Hessberg, P., Pouvesle, N., Winkler, A. K., Schuster, G. and Crowley, J. N.: Phys. Chem. Chem. Phys. 10, 2345-2355, 2008.



Partitioning coefficients ( $K_{ice}$ ) for acetic acid uptake to ice.