

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet V.A2.11 MD11

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>.

This datasheet last evaluated: March 2010; last change in preferred values: March 2010.

HC(O)OH + mineral oxide (dust) surfaces

Experimental data

Parameter	Temp./K	Reference	Technique/ Comments
γ_0			
$(3 \pm 1) \times 10^{-3}$ (dry CaCO_3)	298 *	Al-Hosney et al., 2006	Knudsen-MS (a)
1.7×10^{-5} (Na-Montmorillonite, RH = 0 %)	212	Hatch et al., 2007	Knudsen-MS (b)
1.7×10^{-5} (Na-Montmorillonite, RH = 29 %)			
2.3×10^{-5} (Na-Montmorillonite, RH = 45 %)			

*Experimental temperature not given, presumably room temperature.

Comments

- (a) Continuous flow Knudsen experiments using bulk samples of CaCO_3 prepared from an aqueous slurry by allowing H_2O to evaporate under vacuum overnight. The HC(O)OH concentration was 1.9×10^{11} molecule cm^{-3} . The mass of substrate was varied ($\approx 2 - 70$ mg) to enable a pore diffusion correction to be carried out (all uptake coefficients were derived in this manner). Transmission-FTIR and ATR-FTIR analysis of the substrate were used to obtain mechanistic information.
- (b) Bulk samples of Na-montmorillonite prepared from an aqueous slurry. Concentration of HC(O)OH was varied between $\sim 2-30 \times 10^{11}$ molecule cm^{-3} . The initial uptake coefficient (based on the BET surface area) obtained in first two seconds of exposure was not influenced by slowly changing surface area due to swelling of the clay mineral at non-zero RH and was independent of the formic acid concentration. FTIR sample analysis indicated that the interaction is (at least in part) irreversible.

Preferred Values

none.

Comments on Preferred Values

Al-Hosney et al. (2005) found dominantly (98 %) irreversible uptake of HC(O)OH to calcite. Time and mass dependent uptake coefficients were observed and a pore diffusion correction was performed to derive the “true” uptake coefficient. The tortuosity coefficient (τ) derived from fitting to the mass dependence of the uptake coefficient was 4, slightly larger than the maximum value of 3 suggested to be acceptable for powder samples (Boulter and Marschall, 2006).

Exposure of dry CaCO_3 to $\approx 8 \times 10^{15}$ HC(O)OH resulted in formation of formates and carbonic acid via reaction with $\text{Ca(OH)}(\text{HCO}_3)$ present on CaCO_3 surfaces. Carbonic acid was converted to CO_2 in the presence of moisture, which also results in an enhanced capacity for CaCO_3 to

take up HC(O)OH and an enhanced rate of formate formation. This is explained in terms of surface reorganisation (reactivation) and involvement of underlying layers so that no surface saturation is observed at high relative humidity. Hatch et al. (2007) observed that the initial uptake of formic acid to Na-montmorillonite increased only slightly as RH increased and that the uptake HC(O)OH saturated at calculated coverages that varied between <0.1 and 0.5×10^{14} molecule cm $^{-2}$, which are sub-monolayer. Al Hosney et al derived a saturation surface coverage of $\sim 3 \times 10^{14}$ molecules cm $^{-2}$ for CaCO₃.

The initial uptake coefficients of formic acid to calcite and Na-montmorillonite are vastly different (factor 100), thus uptake to atmospheric dust would be dominated by the calcite fraction. As the uptake of HC(O)OH to calcite was observed to cease at coverages of $\sim 3 \times 10^{14}$ molecules cm $^{-2}$ the initial uptake coefficient is not appropriate for modelling the interaction of HC(O)OH and mineral dust in the atmosphere. Assuming $\sim 3 \times 10^{14}$ reactive sites per cm 2 , and a HC(O)OH mixing ratio of 350 ppt, the initial uptake coefficient would be reduced by a factor of ~ 10 following one hour of exposure. This may be offset by reorganisation of the particle surface at high humidities, but until quantitative kinetic data are available at RH other than zero, we make no recommendation.

References

- Al-Hosney, H. A., Carlos-Cuellar, S., Baltrusaitis, J. and Grassian, V. H.: Phys. Chem. Chem. Phys. 7, 3587-3595, 2005.
Boulter, J. E. and Marschall, J.: J. Phys. Chem. A 110, 10444-10455, 2006.
Hatch, C. D., Gough, R. V. and Tolbert, M. A.: Atmos. Chem. Phys. 7, 4445-4458, 2007.