

Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet HO_x_VOC35

Website: <http://iupac.pole-ether.fr>. See website for latest evaluated data. Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. This data sheet updated: 12th December 2007 (with no revision of the preferred values).

HO + HC(O)OH → products

Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$(4.62 \pm 0.78) \times 10^{-13}$	298-430	Wine et al., 1985	FP/PLP-RF (a)
$(4.90 \pm 0.12) \times 10^{-13}$	296	Jolly et al., 1986	PLP-RA (b)
$2.91 \times 10^{-13} \exp[(102 \pm 194)/T]$	297-445	Singleton et al., 1988	PLP-RA
$(4.47 \pm 0.28) \times 10^{-13}$	297		
$(3.7 \pm 0.4) \times 10^{-13}$	298	Dagaut et al., 1988	FP-RF

Comments

- (a) H atom formation was also measured by resonance fluorescence, and an H-atom formation yield of 0.75 ± 0.25 measured. At 298 K, the measured rate coefficient for the reaction of the HO radical with DC(O)OH was identical to that for HO radical reaction with HC(O)OH.
- (b) Experiments with added O₂ led to non-exponential and slower HO radical decays, indicating the formation of H atoms from the HO radical reaction with HC(O)OH.

Preferred Values

$k = 4.5 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, independent of temperature over the range 290-450 K.

Reliability

$\Delta \log k = \pm 0.15$ at 298 K.


$\Delta(E/R) = \pm 250$ K.

Comments on Preferred Values

A major problem with the determination of the rate coefficient for this reaction concerns the ready dimerization of HC(O)OH. The studies of Wine et al. (1985), Jolly et al. (1986) and Singleton et al. (1988) monitored formic acid in the experimental systems used by UV absorption spectroscopy. The data from these studies (Wine et al., 1985; Jolly et al., 1986; Singleton et al., 1988) agree well, and are in reasonable agreement with the room temperature rate coefficient of Dagaut et al. (1988). The data of Wine et al. (1985) and Singleton et al. (1988) show that the temperature dependence of the rate coefficient is zero within the experimental uncertainties. The average of the rate coefficients of Wine et al. (1985), Jolly et al. (1986) and Singleton et al. (1988) has been used to derive the preferred rate coefficient.

The studies of Wine et al. (1985) and Jolly et al. (1986) showed that H atoms are produced in this reaction, with a yield of 0.75 ± 0.25 (Wine et al., 1985). Furthermore, Wine et al. (1985) and

Singleton et al. (1988) showed that within the experimental uncertainties the rate coefficient for the reaction of the HO radical with DC(O)OH is identical to that for HC(O)OH at 298 K. Also, the room temperature rate coefficients for the reactions of the DO radical with HC(O)OD and DC(O)OD are significantly lower than those for the reactions of the HO radical with HC(O)OH and DC(O)OH (Singleton et al., 1988). The reaction then appears to proceed by,



with overall abstraction of the H (or D) atom from the -OH (or -OD) group being the major pathway at room temperature.

References

- Dagaut, P., Wallington, T. J., Liu, R. and Kurylo, M. J.: *Int. J. Chem. Kinet.*, 20, 331, 1988.
Jolly, G. S., McKenney, D. J., Singleton, D. L., Paraskevopoulos, G. and Bossard, A. R.: *J. Phys. Chem.*, 90, 6557, 1986.
Singleton, D. L., Paraskevopoulos, G., Irwin, R. S., Jolly, G. S. and McKenney, D. J.: *J. Am. Chem. Soc.*, 110, 7786, 1988.
Wine, P. H., Astalos, R. J. and Mauldin III, R. L.: *J. Phys. Chem.*, 89, 2620, 1985.

- Recommendation
- Wine et al. (1985)
- Jolly et al. (1986)
- ▲ Singleton et al. (1988)
- ▼ Dagaut et al. (1988)

