

## IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet V.A5.15 HNNT15

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>.

This data sheet last evaluated: June 2009; last change in preferred values: June 2009.



### Experimental data

Parameter	Temp./K	Reference	Technique/ Comments
$\gamma$			
$3.2 \times 10^{-3}$	200	Hanson and Ravishankara, 1991	CWFT-CIMS (a)

### Comments

- (a) Ice surfaces (7-15  $\mu\text{m}$  thick) were made by vapour deposition and doped with  $\text{HNO}_3$  formed during the  $\text{N}_2\text{O}_5$  uptake (the amount of  $\text{HNO}_3$  was not given). The geometric surface area was used to calculate the uptake coefficient. Experiments were conducted with  $[\text{N}_2\text{O}_5]$  and  $[\text{HCl}]$  at  $\approx 10^9 - 10^{10}$  molecule  $\text{cm}^{-3}$ . HCl was generally in excess, but first order loss of  $\text{N}_2\text{O}_5$  was still observed when  $[\text{N}_2\text{O}_5] / [\text{HCl}] = 2$ . Variation of  $[\text{N}_2\text{O}_5]$  over a factor of 10 did not change the uptake coefficient, a dependence of  $\gamma$  on  $[\text{HCl}]$  was not reported.

### Preferred Values

Parameter	Value	T/K
$\gamma_{\text{gs}}$	$4 \times 10^{-3}$	190 - 210 K
$\theta_{\text{HCl}}$	$7.3 \times 10^{-17} \exp(2858/T) [\text{HCl}] / (7.3 \times 10^{-17} \exp(2858/T) [\text{HCl}] + 1)$	190 - 210 K
<i>Reliability</i>		
$\Delta \log(\gamma)$	0.5	190 - 210 K

### Comments on Preferred Value


The single study (Hanson and Ravishankara, 1991) of the reaction of  $\text{N}_2\text{O}_5$  and HCl on a NAT surface shows that the  $\text{N}_2\text{O}_5$  uptake proceeds more efficiently than on NAT alone (for which a value of  $\gamma = 6 \times 10^{-4}$  was obtained), indicating that a surface reaction takes place presumably to form  $\text{HNO}_3$  and  $\text{ClNO}_2$ , though these products were not seen. Application of the reported value of  $\gamma$  to the atmosphere requires parameterisation of the HCl surface coverage on NAT. We assume that this is the same as for pure ice to derive the expression for  $\theta_{\text{HCl}}$  (see HCl + ice datasheet), which can be combined with a value of  $\gamma_{\text{gs}}$  of  $4 \times 10^{-3}$  to give net uptake coefficients in line with those observed by Hanson and Ravishankara, 1991.

$$\gamma = \gamma_{\text{gs}} \theta_{\text{HCl}} + 6 \times 10^{-4}$$

This expression assumes that the reaction is driven by the availability of surface HCl and the temperature dependence is controlled only by HCl coverage. As  $\text{HCl} \rightarrow 0$ , the uptake coefficient should approach that on pure NAT (i.e.  $6 \times 10^{-4}$ ). The ratio of  $\text{ClNO}_2$  formation to the overall uptake is given by  $(\gamma - 6 \times 10^{-4}) / \gamma$ .

### References

Hanson, D. R. and Ravishankara, A. R.: J. Geophys. Res. 96, 5081-5090, 1991.



Uptake coefficients for  $\text{N}_2\text{O}_5$  reaction on a HCl doped NAT surface according to the expression given in the datasheet. The single experimental dataset is consistent with this parameterisation, but does not confirm it.