


## Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet NO<sub>3</sub>\_VOC38

Datasheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>)

This datasheet last evaluated: Nov. 2016; last change in preferred values: June 2013



### Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$(1.07 \pm 0.16) \times 10^{-11}$ $2.3 \times 10^{-10} \exp[-(940 \pm 200)/T]$	298 298-393	Martínez et al., 1999	DF-LIF (a)
<i>Relative Rate Coefficients</i>			
$(1.01 \pm 0.03) \times 10^{-11}$	296	Atkinson et al., 1990	RR (b)

Sabinene is 1-isopropyl-4-methylene-bicyclo[3.1.0]hexane.

### Comments

- (a) NO<sub>3</sub> radicals ( $6\text{-}30 \times 10^{11} \text{ molecule cm}^{-3}$ ) generated from reaction of F atoms (made in a microwave discharge through F<sub>2</sub>/He) with HNO<sub>3</sub>. Flow tube was operated at ~1.33 mbar (1 Torr) He at 4 temperatures between 298 and 393 K. Sabinene was present at similar concentrations (1-3 fold) to NO<sub>3</sub>. So that absolute NO<sub>3</sub> concentrations (derived by titration with tetramethylethene) were necessary to derive the rate coefficient.
- (b) Relative rate of loss of sabinene and 2-methyl-2-butene (reference reactant) in a 6400 L Teflon chamber at 980 mbar (735 Torr) of air was monitored by GC. NO<sub>3</sub> was generated by the thermal decomposition of N<sub>2</sub>O<sub>5</sub>. The rate constant ratio,  $k(\text{NO}_3 + \text{sabinene}) / k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 1.08 \pm 0.03$  is placed on an absolute basis using  $k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 9.37 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K (Atkinson and Arey, 2003).

### Preferred Values

Parameter	Value	T/K
$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$1.0 \times 10^{-11}$	298
<i>Reliability</i>		
$\Delta \log k$	$\pm 0.10$	298

## Comments on Preferred Values


The preferred value at 298 K is based on the relative rate study of Atkinson et al. (1990). The significant, positive dependence of  $k$  on temperature observed by Martínez et al. (1999) requires validation.

There are no product studies of this reaction, though the large rate constant indicates that the reaction proceeds mainly via addition of  $\text{NO}_3$  across a double bond to form a chemically activated nitro-oxyalkyl radical. At pressures found in the troposphere this adduct will undergo collisional stabilization prior to reaction with  $\text{O}_2$  to form a nitrooxyalkyl peroxy radical.

Secondary organic aerosol has been observed in the smog-chamber studies of  $\text{NO}_3$  + sabinene (Fry et al., 2014) with mass-based yields of up to 45 % (at  $10 \mu\text{g m}^{-3}$  aerosol loading).

## References

- Atkinson, R., Aschmann, S. M., and Arey, J., *Atmos. Env. A*, 24, 2647-2654, 1990  
Atkinson, R., and Arey, J., *Chem. Rev.*, 103, 4605-4638, 2003.  
IUPAC: Task Group on Atmospheric Chemical Kinetic Data Evaluation, (Ammann, M., Cox, R.A., Crowley, J.N., Herrmann, H., Jenkin, M.E., McNeill, V.F., Mellouki, A., Rossi, M. J., Troe, J. and Wallington, T. J.) <http://iupac.pole-ether.fr/index.html>, <http://iupac.pole-ether.fr/index.html>, 2016.  
Martínez, E., Cabañas, B., Aranda, A., Martín, P., and Salgado, S., *J. Atmos. Chem.*, 33, 265-282, 1999.



Rate coefficients for  $\text{NO}_3$  + sabinene