


Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet NO3_VOC48

Datasheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>)

This datasheet last evaluated: Nov. 2016; last change in preferred values: Nov. 2016



Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Relative Rate Coefficients</i>			
$(8.16 \pm 0.72) \times 10^{-12}$	296	Atkinson et al., 1990	RR (a)

α -cedrene is: (1*S*,2*R*,5*S*,7*R*)-2,6,6,8-tetramethyltricyclo[5.3.1.0^{1,5}]undec-8-ene

Comments

- (a) Relative rate of loss of α -cedrene and 2-methyl-2-butene (reference reactant) in a 6700 L Teflon chamber at 987 mbar (740 Torr) of air was monitored by GC-FID. NO_3 was generated by the thermal decomposition of N_2O_5 . The rate constant ratio, $k(\text{NO}_3 + \alpha\text{-cedrene}) / k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 0.871 \pm 0.077$ is placed on an absolute basis using $k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 9.37 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298 K (Atkinson and Arey, 2003).

Preferred Values

Parameter	Value	T/K
$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	8.2×10^{-12}	298
<i>Reliability</i>		
$\Delta \log k$	± 0.25	298

Comments on Preferred Values

The preferred value at 298 K is based on the relative rate study of Shu and Atkinson (1995). The error limits have been expanded to reflect the fact that this is the only study available. The rate coefficient is consistent with estimates based on the number of double bonds and the number and position of the alkyl substituents around them (Atkinson 1991).

There are no studies of the gas-phase products of this reaction, though it is expected to proceed predominantly via addition of NO_3 across the double bond to form a nitrooxyalkyl radical which can react with O_2 to form a nitrooxyalkyl peroxy radical.

Secondary aerosol is formed efficiently in chamber studies of this reaction (Jaoui et al., 2013).

References

- Atkinson, R., *J. Phys. Chem. Ref. Data*, 20, 459-507, 1991.
Atkinson, R., and Arey, J., *Chem. Rev.*, 103, 4605-4638, 2003.
Jaoui, M., Kleindienst, T. E., Docherty, K. S., Lewandowski, M., and Offenberg, J. H.,
Environmental Chemistry, 10, 178-193, 2013.
Shu, Y. H., and Atkinson, R., *J. Geophys. Res. -Atmos.*, 100, 7275-7281, 1995.