

# IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet I.A3.33 NO<sub>x</sub>3

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>).

This data sheet last evaluated: 4<sup>th</sup> June 2009; last change in preferred values April 2005.



$$\Delta H^\circ = -208.6 \text{ kJ}\cdot\text{mol}^{-1}$$

## Low-pressure rate coefficients Rate coefficient data

$k_0/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$3 \times 10^{-31} (T/300)^{-1.75} [\text{N}_2]$	220-296	Burkholder and Ravishankara, 2000	PLP (a)
$(1.3 \pm 0.3) \times 10^{-31} (T/300)^{-1.5} [\text{N}_2]$	300-400	Hahn et al., 2000	PLP (b)
<i>Relative Rate Coefficients</i>			
$(9.2 \pm 1) \times 10^{-32} [\text{N}_2]$	297	Harker and Johnston, 1973	RR (c)
$(8.0 \pm 1) \times 10^{-32} [\text{N}_2]$	295	Hippler et al., 1975	RR (d)

## Comments

- (a) Oxygen atoms generated by laser photolysis of NO<sub>2</sub> at 352 nm, NO<sub>3</sub> monitored by long-path diode laser absorption at 662 nm. Measurements were made in N<sub>2</sub> over the pressure range 20-800 Torr; falloff curves were represented with  $F_c = 0.6$  and  $k_\infty = 3.75 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  independent of the temperature.
- (b) Oxygen atoms generated by laser photolysis of N<sub>2</sub>O at 193 nm, NO<sub>3</sub> monitored by light absorption at 578 nm. Falloff curves measured at 300 and 400 K in N<sub>2</sub> over the pressure range 1-900 bar. Falloff curves represented with  $F_c = 0.71 \exp(-T/1700)$ , which corresponds to  $F_c = 0.6$  at 300 K, and  $k_\infty = (2.3 \pm 0.2) \times 10^{-11} (T/300)^{0.24} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .
- (c) O(<sup>3</sup>P) atoms were generated by the photolysis of NO<sub>2</sub> in the presence of 1 bar of N<sub>2</sub>. NO<sub>2</sub> and N<sub>2</sub>O<sub>5</sub> concentrations were monitored by IR absorption. The measured value of  $k/k(\text{O} + \text{NO}_2)$  was evaluated with  $k(\text{O} + \text{NO}_2) = 9.3 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ . The rate coefficient has been reevaluated by increasing the measured rate coefficient by 10% to account for a 10% falloff below  $k_0$ , as measured by in ref. 5.
- (d) O(<sup>3</sup>P) atoms were generated by the photolysis of NO<sub>2</sub> at various N<sub>2</sub> pressures. NO<sub>2</sub> was monitored by UV absorption. The measured value of  $k/k(\text{O} + \text{NO}_2)$  was evaluated with  $k(\text{O} + \text{NO}_2) = 9.3 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , taking N<sub>2</sub>O<sub>5</sub> reactions and falloff effects into account.

## Preferred Values

$$k_0 = 1.3 \times 10^{-31} (T/300)^{-1.5} [\text{N}_2] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ over the temperature range 200-400 K.}$$

## Reliability

$$\Delta \log k_0 = \pm 0.30 \text{ at 298 K.}$$

$$\Delta n = \pm 1.$$

### Comments on Preferred Values

The preferred values are from the falloff extrapolations of the absolute rate measurements of Hahn et al. (2000) and the relative rate measurements of Harker and Johnston (1973), Hippler et al. (1975), Gaedtke and Troe (1975), Schuck et al. (1966) and Troe (1969). The discrepancy between the absolute rate measurements below 800 Torr from ref. 1 and above from Hahn et al. (2000) may be due to difficulties in separating the reactions  $O + NO_2 \rightarrow O_2 + NO$  and  $O + NO_2 + M \rightarrow NO_3 + M$  with subsequent  $NO_2 + NO_3 + M \rightarrow N_2O_5 + M$ . Because the data from Hahn et al. (2000) are in much better consistency with the relative rate measurements from Hahn et al. (2000), they are preferred here. It should be emphasized that falloff effects are important at atmospheric pressures; these effects are represented with  $F_c = 0.6$  and  $k_\infty = 2.3 \times 10^{-11} (T/300)^{0.24} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  from the measurements and theoretical analysis of Hahn et al. (2000).

### High-pressure rate coefficients Rate coefficient data

$k_\infty/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i> $(2.3 \pm 0.2) \times 10^{-11} (T/300)^{0.24}$	300-400	Hahn et al., 2000	PLP (a)
<i>Relative Rate Coefficients</i> $(2.2 \pm 0.4) \times 10^{-11}$	300	Gaedtke and Troe, 1975	(b)

### Comments

- (a) See comment (b) for  $k_0$ .  
 (b) Analysis of the pressure dependence of the quantum yield of  $NO_2$  photolysis between 1 and 1000 bar of  $N_2$ . Measurements of  $k_\infty/k(O + NO_2 \rightarrow O_2 + NO)$  evaluated with  $k(O + NO_2) = 9 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .

### Preferred Values

$k_\infty = 2.3 \times 10^{-11} (T/300)^{0.24} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  over the temperature range 200-400 K.

### Reliability

$\Delta \log k_\infty = \pm 0.2$  over the temperature range 200-400 K.

### Comments on Preferred Values

The absolute and relative rate measurements of the falloff curve over a very wide pressure range from Hahn et al. (2000) and Gaedtke and Troe (1975), on which the preferred values are based, allow for a reliable extrapolation to the high pressure limit of the reaction. Falloff curves are constructed with  $F_c = 0.6$  and  $k_0 = 1.3 \times 10^{-31} (T/300)^{-1.5} [N_2] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .

The following text-line combines the preferred values for the high and low pressure limiting rate coefficients to generate a single, cut-and-paste expression for calculation of  $k$ :

$$= ((1.3e-31*(T/300)^{-1.5}*M*(2.3e-11*(T/300)^{0.24})/((1.3e-31*(T/300)^{-1.5}*M+(2.3e-11*(T/300)^{0.24}))*10^{(\log10(0.6)/(1+(\log10((1.3e-31*(T/300)^{-1.5}*M/(2.3E-11*(T/300)^{0.24}))/((0.75-1.27*\log10(0.6))))^2))$$

The molecular density,  $M = 7.243 \times 10^{21} P(\text{bar})/T(\text{K})$

## References

- Burkholder, J. B. and Ravishankara, A. R.: J. Phys. Chem. A 104, 6752, 2000.  
Gaedtke, H. and Troe, J.: Ber. Bunsenges. Phys. Chem. 79, 184, 1975.  
Hahn, J., Luther, K. and Troe, J.: Phys. Chem. Chem. Phys. 2, 5098, 2000.  
Harker, A. B. and Johnston, H. S.: J. Phys. Chem. 77, 1153, 1973.  
Hippler, H., Schippert, C. and Troe, J.: Int. J. Chem. Kinet. Symp. 1, 27, 1975.  
Schuck, E. A., Stephens, E. R. and Schrock, R. R.: J. Air Pollut. Contr. Assoc. 16, 695, 1966.  
Troe, J.: Ber. Bunsenges. Phys. Chem. 73, 906, 1969.