

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet Ox_AROM4

Website: <http://iupac.pole-ether.fr>. See website for latest evaluated data. Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission.

This data sheet last evaluated September 2008; last change in preferred values September 2008.

O₃ + *o*-CH₃C₆H₄OH (*o*-cresol) → products

Rate coefficient data

<i>k</i> /cm ³ molecule ⁻¹ s ⁻¹	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
~6 × 10 ⁻¹⁹	300 ± 1	Atkinson et al., 1978	S-GC (a)
(2.55 ± 0.39) × 10 ⁻¹⁹	296 ± 2	Atkinson et al., 1982	S-CL (b)

Comments

- (a) *o*-Cresol disappearance monitored by GC in the presence of excess O₃. Experiments were carried out in a ~5500 L Teflon chamber at atmospheric pressure of air. The cited rate was stated to be uncertain by a factor of ~2 and may be an upper limit due to the possibility of secondary reactions removing *o*-cresol.
- (b) O₃ decays were monitored by chemiluminescence in the presence of excess *o*-cresol in a ~175 L Teflon chamber at atmospheric pressure of air.

Preferred Values

$$k = 2.6 \times 10^{-19} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \text{ at } 298 \text{ K.}$$

Reliability

$$\Delta \log k = \pm 0.3 \text{ at } 298 \text{ K.}$$

Comments on Preferred Values

The preferred value is based on the study of Atkinson et al. (1982) in which O₃ decays were monitored in the presence of excess *o*-cresol. The approximate rate coefficient of Atkinson et al. (1978), which may be an upper limit due to secondary reactions removing *o*-cresol, is reasonably consistent with that of Atkinson et al. (1982).

References

- Atkinson, R., Aschmann, S. M., Fitz, D. R., Winer, A. M. and Pitts Jr., J. N.: Int. J. Chem. Kinet., 14, 13, 1982.
- Atkinson, R., Darnall, K. R. and Pitts Jr., J. N.: J. Phys. Chem., 82, 2759, 1978.