

Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet Ox_VOC38

Datasheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission.

The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>)

This datasheet last evaluated: August 2018; last change in preferred values: June 2015



Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$(8.8 \pm 0.8) \times 10^{-18}$	295 ± 2	Richters et al., 2015	S-IR/UVA (a)
<i>Relative Rate Coefficients</i>			
$(1.1 \pm 0.5) \times 10^{-15}$	366 ± 2	Ghalaieny et al., 2012	RR-GC (b)
$(1.57 \pm 0.29) \times 10^{-17}$	295 ± 2	Richters et al., 2015	RR-MS (c)
$(7.56 \pm 0.21) \times 10^{-18}$	295 ± 2	Richters et al., 2015	RR-MS (d)

isolongifolene is (1R)-2,2,7,7-tetramethyltricyclo[6.2.1.0^{1,6}]undec-5-ene

Comments

- k determined from the observed pseudo-first order rate of ozone decay (measured by UVA at 254 nm) in the presence of known excess concentrations of isolongifolene (measured by FTIR), in stopped-flow experiments at a total pressure of ~ 1 bar, with sufficient propane to scavenge $>99\%$ of HO radicals.
- The concentrations of isolongifolene and 2,3-dimethyl-but-2-ene (the reference compound), with excess cyclohexane to scavenge HO radicals, were monitored by GC-FID in a 123 L Teflon-coated chamber at 780 Torr (1040 mbar) pressure of N_2 , with repeated injections of O_3/O_2 . The measured rate coefficient ratio, $k(\text{O}_3 + \text{isolongifolene})/k(\text{O}_3 + 2,3\text{-dimethyl-but-2-ene}) = 0.87$, is placed on an absolute basis using $k(\text{O}_3 + 2,3\text{-dimethyl-but-2-ene}) = 1.32 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 366 K (IUPAC, current recommendation). It is noted that the authors used a much lower value of $k(\text{O}_3 + 2,3\text{-dimethyl-but-2-ene}) = 2.89 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, based on a relative rate measurement reported in the same study, leading to a reported value of $k = (2.5 \pm 1.1) \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 366 K.
- The concentrations of isolongifolene and 2-methyl-but-2-ene (the reference compound), with propane to scavenge HO radicals, were monitored by PTR-MS in a flow tube at atmospheric pressure. The measured rate coefficient ratio, $k(\text{O}_3 + \text{isolongifolene})/k(\text{O}_3 + 2\text{-methyl-but-2-ene}) = (0.040 \pm 0.004)$, is placed on an absolute basis using $k(\text{O}_3 + 2\text{-methyl-but-2-ene}) = 3.92 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 295 K (Atkinson and Arey, 2003).
- The concentrations of isolongifolene and *cis*-but-2-ene (the reference compound), with propane to scavenge HO radicals, were monitored by PTR-MS in a flow tube at atmospheric pressure. The

measured rate coefficient ratio, $k(\text{O}_3 + \text{isolongifolene})/k(\text{O}_3 + \text{cis-but-2-ene}) = (0.060 \pm 0.004)$, is placed on an absolute basis using $k(\text{O}_3 + \text{cis-but-2-ene}) = 1.26 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 295 K (IUPAC, current recommendation).

Preferred Values

Parameter	Value	T/K
$k / \text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	1.0×10^{-17}	298
<i>Reliability</i>		
$\Delta \log k$	± 0.3	298

Comments on Preferred Values

The preferred value of the rate coefficient at 298 K is based on a rounded average of the absolute and relative rate determinations of Richters et al. (2015), with an uncertainty that encompasses the range of values. Although Ghalaieny et al. (2012) report a comparable value of k at 366 K (see comment (c)), this was apparently based on use of an anomalously low value of the rate coefficient for the reference reaction of O_3 with 2,3-dimethyl-but-2-ene. The value tabulated above is based on the current IUPAC recommendation (see comment (c)), leading to a value of k (at 366 K) that is two orders of magnitude higher than the preferred value.

The reaction is expected to proceed via addition of O_3 to the C=C bond, with the subsequent mechanism following the general framework outlined in the datasheets for other mono- and sesquiterpenes possessing endocyclic C=C bonds (e.g. α -cedrene). However, there have apparently been no product or mechanistic studies to confirm this.

References

- Atkinson, R. and Arey, J.: Chem. Rev., 103, 4605, 2003.
Ghalaieny, M., Bacak, A., McGillen, M., Martin, D., Knights, A. V., O'Doherty, S., Shallcross, D. E. and Percival, C. J.: Phys. Chem. Chem. Phys., 14, 6596, 2012.
Richters, S., Herrmann, H. and Berndt, T.: Phys. Chem. Chem. Phys., 17, 11658, 2015.