

## Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet V.A5.16 HNNT16

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>. This data sheet last evaluated: January 2009; last change in preferred values: January 2009.



### Experimental data

Parameter	Temp./K	Reference	Technique/ Comments
$\gamma$			
$5 \times 10^{-3}$	200	Hanson and Ravishankara, 1992	CWFT-CIMS (a)

### Comments

- (a) Ice surfaces (2-10  $\mu\text{m}$  thick) were made by vapour deposition and doped with  $\text{HNO}_3$  (the amount of  $\text{HNO}_3$  was not given). The geometric surface area was used to calculate the uptake coefficient. Experiments were conducted with  $5 \times 10^9 - 10^{11}$  molecule  $\text{cm}^{-3}$  HBr and  $10^{11} - 10^{12}$  molecule  $\text{cm}^{-3}$   $\text{N}_2\text{O}_5$ . Values of the uptake coefficient varied between a high value of  $\approx 4 \times 10^{-2}$  and a low value of  $5 \times 10^{-3}$ , the latter being obtained at the lower HBr concentrations.

### Preferred Values

Parameter	Value	T/K
$\gamma_{\text{ER}}$	$2 \times 10^{-2}$	
$\theta_{\text{HBr}}$	$4.14 \times 10^{-10} [\text{HBr}]^{0.88}$	
<i>Reliability</i>		
$\Delta \log \gamma$	0.5	200

### Comments on Preferred Values

There is a single study (Hanson and Ravishankara, 1992) of the reaction of  $\text{N}_2\text{O}_5$  and HBr on a NAT-like (or  $\text{HNO}_3$  doped) surface. The uptake coefficient was found to be enhanced compared to  $\text{N}_2\text{O}_5$  uptake to pure NAT ( $\gamma \leq 1 \times 10^{-3}$ ), indicative of surface reaction with possible products  $\text{BrONO}$  and  $\text{HNO}_3$  (not observed). In order to parameterise the uptake coefficient,  $\gamma$ , we have assumed an Eley-Rideal type mechanism with the surface coverage of HBr the same as that for pure ice.

$\gamma = \gamma_{\text{gs}} \theta_{\text{HBr}}$ , with  $[\text{HBr}]$  in molecule  $\text{cm}^{-3}$ .

The parameterisation above yields a value of  $\gamma_{\text{net}} = 5 \times 10^{-3}$  at concentrations of HBr close to  $10^{10}$  molecule  $\text{cm}^{-3}$ , increasing to  $\gamma = 4 \times 10^{-2}$  at HBr close to  $10^{11}$  molecule  $\text{cm}^{-3}$ , which are consistent with the experimental observations.

## References

Hanson, D. R. and Ravishankara, A. R.: J. Phys. Chem. 96, 9441-9446, 1992.

