

## Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet V.A5.13 HNNT13

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>. This data sheet last evaluated: May 2008; last change in preferred values: May 2008.



### Uptake coefficient data

Parameter	Temp/K	Reference	Comment
<i>Uptake Coefficients( ClONO<sub>2</sub>)</i>			
0.3(+0.7, -0.1)		Hanson and Ravishankara, 1991	CWFT-CIMS(a)
0.27±0.04	196	Leu, Moore and Keyser, 1991	CWFT-MS(b)
0.3(+0.7, -0.1)	191	Hanson and Ravishankara, 1992	CWFT-CIMS(c)
$\gamma_{ss} \geq 0.20$ 100% RH	202	Abbatt and Molina, 1992	CWFT-MS(d)
$\gamma_{ss} = (3.0 \pm 1.0) \times 10^{-3}$ 25% RH			
$\gamma_{ss} = 0.23$ (p <sub>HCl</sub> =6.7x10 <sup>-8</sup> mbar)	190	Hanson and Ravishankara, 1993	CWFT-CIMS(e)
90% RH	190		
$\gamma_{ss} = 0.20$ (p <sub>HCl</sub> ~4.6x10 <sup>-7</sup> mbar)	197		
90%RH			
$\gamma_{ss} = 0.03$ (p <sub>HCl</sub> ~4.6x10 <sup>-7</sup> mbar)			
30%RH			

### Comments

- (a) Vapour deposited ice; NAT was prepared in situ by converting N<sub>2</sub>O<sub>5</sub> into HNO<sub>3</sub> on the ice surface well past saturation. The HNO<sub>3</sub> vapor detected at the downstream end of the flow tube was consistent within a factor of two with the expected vapor pressure over NAT/ HNO<sub>3</sub> in ice solid solution near 201K. [HCl]<sub>0</sub> = 2 x [ClONO<sub>2</sub>]<sub>0</sub>, i.e. ~ 0.4-1.2 x 10<sup>10</sup> molecule cm<sup>-3</sup>.
- (b) The films, typically 70 mμ thick, were prepared in situ by vapour condensation of HNO<sub>3</sub> and H<sub>2</sub>O at 196K. Initial p<sub>HCl</sub> = 0.27 to 2.7x10<sup>-3</sup> mbar, P<sub>ClONO<sub>2</sub></sub>=10.6 x10<sup>-5</sup> mbar. The measured values of γ were independent of the composition of HNO<sub>3</sub>/NAT in the range 41.8 to 60.4% and of the HCl content (0.0375% to 3.91%HCl). The authors give a corrected value of γ =0.10±0.02 if pore diffusion is taken into account.

- (c) Details under (a). The uptake of ClONO<sub>2</sub>, HCl and the formation of the reaction product Cl<sub>2</sub> were studied at constant HCl concentration (10<sup>10</sup> molecule cm<sup>-3</sup>) and varying ClONO<sub>2</sub> concentration ranging from 0.6-3.0 x 10<sup>10</sup> molecule cm<sup>-3</sup>. The authors argue in favor of a direct reaction between ClONO<sub>2</sub> and HCl rather than a reaction via the intermediate HOCl in view of the high value for  $\gamma$ .
- (d) The NAT films were prepared starting from 10 mμ thick ice films exposed to small pressures of HNO<sub>3</sub> over long periods of time resulting in a 0.1 mμ thick NAT layer on top of the ice film. pressure of ClONO<sub>2</sub> ranging from 1.3 to 12 x 10<sup>-6</sup> mbar and HCl ranging from 5.3 to 13.3x10<sup>-6</sup> mbar. A factor of four increase in  $\gamma$  (from 0.01 to 0.04) was noted when p<sub>HCl</sub> increased from 2 to 10.6 x 10<sup>-6</sup> mbar.
- (e) Details under (a and c). A 0.05 mm thick NAT film was grown on a 0.5 mm thick H<sub>2</sub>O ice undercoat by flowing HNO<sub>3</sub> at 1.3x10<sup>-6</sup> mbar. Subsequently the ice undercoat was evaporated. p(H<sub>2</sub>O) added to He flow to adjust relative humidity over the film.  $\gamma$  decreased strongly with decreasing relative humidity which was varied by adjusting the temperature of the frozen phase at a constant H<sub>2</sub>O flow rate of 3.3x10<sup>-4</sup> mbar.  $\gamma$  given by the expression:  $1/\gamma = 1/\gamma_{\max} + 1/A\exp(b\Delta T)$  where  $\Delta T$  is (T-190), the difference between the temperature of interest and the ice point temperature at which RH is 100% at the chosen flow rate of H<sub>2</sub>O (RH=100 x p<sub>H2O</sub>/p<sub>ice</sub>(T)). For p<sub>HCl</sub> ≈ 6.7x10<sup>-8</sup> mbar and ≈ 4.6 x10<sup>-7</sup> mbar ([HCl] (2.5 and 17.5 x 10<sup>9</sup> molecule cm<sup>-3</sup> at 195 K),  $\gamma_{\max}$  = 0.23 and 0.20, A=0.7022 and 2.2543, b = -0.518 and -0.558 respectively.

### Preferred values

Parameter	Value		Temp/K
$\gamma_{\text{rxn}}$	$[1/ \gamma_{\text{max}} + 1/A\exp(B\Delta T)]^{-1}$		190 -200
pH <sub>2</sub> O (mbar )	3.3 x 10 <sup>-4</sup>		(RH=20-100%)
$\Delta T$ (K)	T - 190		
pHCl (mbar )	6.7x10 <sup>-8</sup>	4.6 x10 <sup>-7</sup>	
$\gamma_{\text{max}}$	0.25	0.20	
A	0.7022	2.2543	
B	-0.518	-0.558	
<i>Reliability</i>			
$\Delta \log( \gamma_{\text{rxn}} )$	±0.3		190 – 200 K.

### Comments on Preferred Values

As with ice films the uptake of ClONO<sub>2</sub> on NAT films in the presence of HCl is followed by reaction to form Cl<sub>2</sub> and HNO<sub>3</sub> in a surface reaction. At stratospheric temperatures Cl<sub>2</sub> partitions into the gas phase, but HNO<sub>3</sub> remains at the surface with formation of hydrates. The uptake coefficients of ClONO<sub>2</sub> in the presence of HCl on H<sub>2</sub>O-rich NAT substrates are similar to those on pure ice, but show a strong dependence on relative humidity. Thus  $\gamma$  decreases with decreasing p(H<sub>2</sub>O) and decreases with increasing temperature at fixed p(H<sub>2</sub>O).

This reflects the amounts of surface-adsorbed water and reactants, and leads to a complex dependence of  $\gamma$  with conditions.

Uptake coefficients measured on water-rich NAT (100% rh) from the different studies agree quite well. At lower RH there is more variability. Only Hanson and Ravishankara and Abbatt and Molina did a systematic study of the water dependence; in the former study RH was varied by changing T at constant  $p(\text{H}_2\text{O})$  and they observed less dependence of  $\gamma$  compared to Abbatt and Molina who varied  $p(\text{H}_2\text{O})$  at constant T; the latter used higher reactant concentrations which could have led to more influence of  $\text{HNO}_3$  product, reducing surface water availability. On  $\text{H}_2\text{O}$ -rich NAT substrates Hanson and Ravishankara observed that  $\gamma$  increases slowly with  $p_{\text{HCl}}$  in the range  $\approx (0.5-5) \times 10^{-7}$  mbar, which is consistent with the high fractional surface coverage for HCl ( $\theta \approx 0.33$ ) for these conditions. At lower RH and higher [HCl], Abbatt and Molina observed much stronger dependence of  $p_{\text{HCl}}$  in the range  $\approx (2-10) \times 10^{-6}$

The preferred values for the reactive uptake coefficient on NAT under stratospheric conditions are given by the parameterisation of Hanson & Ravishankara (1993), which gives the reactive uptake coefficient for specified [HCl] and fixed  $p(\text{H}_2\text{O}) = 3.3 \times 10^{-4}$  mbar, as a function of temperature in the range 190 -200 K. This corresponds to RH in the range 20-100%.

In view of the complex dependence of the uptake coefficient on the state of the  $\text{HNO}_3$ -rich surfaces, and the lack of consistency in the reported data for these conditions, no recommendation is made for  $\gamma$  at lower [HCl] and absolute humidity. For uptake on NAT surfaces with at lower [HCl] and absolute humidity in the NAT stability region, a parameterisation for  $\gamma$  using a Langmuir-Hinshelwood model such as used for  $\text{ClONO}_2 + \text{HCl}$  on ice (IUPAC, 2009), would require a better definition of the surface concentration of water and HCl on NAT than is available at present.

## References

- Abbatt, J.P.D. and Molina, M.J.: J. Phys. Chem. 96, 7674 (1992).
- Hanson, D.R. and Ravishankara, A.R.: J. Geophys. Res. 96, 5081 (1991).
- Hanson, D.R. and Ravishankara, A.R.: J. Geophys. Res. 98, 22931 (1993).
- Hanson, D.R. and Ravishankara, A.R.: J. Phys. Chem. 96, 2682 (1992).
- Leu, M.-T., Moore, S.B. and Keyser, L.F.: J. Phys. Chem. 95, 7763 (1991).

Figure 1. The dependence of the steady-state uptake coefficient of  $\text{ClONO}_2$  on NAT surfaces as a function of temperature range 190 – 200 K. Prediction for  $p(\text{HCl}) = 6.7 \times 10^{-8}$  mbar and  $\approx 4.6 \times 10^{-7}$  mbar

